



United Behavioral Health

Supplemental Clinical Criteria: Applied Behavior Analysis

Document Number: BH803ABA032020

Effective Date: March 16, 2020

Table of Contents

[Introduction](#)

[Instructions for Use](#)

[Benefit Considerations](#)

[State Mandates](#)

[Description of Service](#)

[Coverage Rationale](#)

[Applicable Codes](#)

[References](#)

[Revision History](#)

INTRODUCTION

Supplemental Clinical Criteria are a set of objective and evidence-based behavioral health criteria used by medical necessity plans to standardize coverage determinations, promote evidence-based practices, and support members' recovery, resiliency, and wellbeing for behavioral health benefit plans that are managed by Optum®¹.

INSTRUCTIONS FOR USE

This guideline is used to make coverage determinations as well as to inform discussions about evidence-based practices and discharge planning for behavioral health benefit plans managed by Optum. When deciding coverage, the member's specific benefits must be referenced.

All reviewers must first identify member eligibility, the member-specific benefit plan coverage, and any federal or state regulatory requirements that supersede the member's benefits prior to using this guideline. In the event that the requested service or procedure is limited or excluded from the benefit, is defined differently or there is otherwise a conflict between this guideline and the member's specific benefit, the member's specific benefit supersedes this guideline. Other clinical criteria may apply. Optum reserves the right, in its sole discretion, to modify its clinical criteria as necessary using the process described in *Clinical Criteria*.

This guideline is provided for informational purposes. It does not constitute medical advice.

Optum may also use tools developed by third parties that are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

Optum may develop clinical criteria or adopt externally-developed clinical criteria that supersede this guideline when required to do so by contract or regulation.

BENEFIT CONSIDERATIONS

Before using this policy, please check the member-specific benefit plan document and any federal or state mandates, if applicable.

¹ Optum is a brand used by United Behavioral Health and its affiliates.

STATE MANDATES

For Arizona Medicaid Developmental Disabilities members, Applied Behavior Analysis (ABA) is covered for the treatment of developmental disorders not limited to Autism Spectrum Disorder when all other criteria are met.

For Indiana members refer to the following:

Services are intensive and may be provided daily. All determinations must be based on the individualized objectives of the treatment plan and unique needs of the member. No quantitative benefit coverage limitations are implied by reference to these guidelines. The intensity of service should consider the member's ability to participate, benefit, and tolerate the full spectrum of services received concurrently, including non-behavioral services such as school, speech and occupational therapies. Treatment intensity should be increased or decreased based on the member's response and current needs.

Refer to the following for fully insured policies in Maryland

Use the following criteria as specified in the Code of Maryland Regulations (*MD COMAR 31.10.39.03. April 3, 2014*):

- A. Applied Behavioral Analysis (ABA) services include behavioral health treatment, psychological care, and therapeutic care of members diagnosed Autism Spectrum Disorder.
- B. The following are required for the initiation and continuation of ABA services:
 1. A comprehensive evaluation of a child by the child's primary care provider or specialty physician identifying the need for treatment of Autism Spectrum Disorder.
 2. A prescription from a child's primary care provider or specialty physician that includes specific treatment goals.
 3. Annual review by the prescribing primary care provider or specialty physician, in consultation with the ABA provider, that includes:
 - i. Documentation of benefit to the child;
 - ii. Identification of new or continuing treatment goals; and
 - iii. Development of a new or continuing treatment plan.
- C. Applied Behavioral Analysis (ABA) Services that meet the above criteria will not be denied solely on the number of hours of habilitative services prescribed for:
 1. Up to 25 hours per week for members between the ages of 18 months and 5 years old.
 2. Up to 10 hours per week for members between the ages of 6 and 18 years old
 3. Additional hours of ABA services will be authorized if determined to be medically necessary and appropriate.
 4. Services are only delivered by providers who are licensed, certified, or otherwise authorized under the Health Occupations Article or similar licensing, certification, and authorization requirements of another state or U.S. territory where ABA services are provided.
- D. Location of Services
 1. ABA services are not denied if the treatment plan identifies the child's school as the location of services.
 2. Services are not authorized under an Individualized Education Program (IEP) or any obligation imposed on a public school by the Individuals with Disabilities Education Act.
- E. ABA will not be denied on the basis that it is experimental or investigational.

DESCRIPTION OF SERVICE

Applied Behavior Analysis

ABA is a scientific discipline among the helping professions that focuses on the analysis, design, implementation, and evaluation of social and other environmental modifications to produce meaningful changes in human behavior. ABA includes the use of direct observation, measurement, and functional analysis of the relations between environment and behavior. ABA uses changes in environmental

events, including antecedent stimuli and consequences, to produce practical and significant changes in behavior. These relevant environmental events are usually identified through a variety of specialized assessment methods. ABA is based on the fact that an individual's behavior is determined by past and current environmental events in conjunction with organic variables such as their genetic endowment and physiological variables. Thus, when applied to ASD, ABA focuses on treating the problems of the disorder by altering the individual's social and learning environments. (Behavior Analyst Certification Board Inc., 2014).

COVERAGE RATIONALE

Applied Behavior Analysis (ABA) is proven for the treatment of autism spectrum disorder in children when the following conditions are met:

- The intervention is a systematic approach, based on the principles of comprehensive applied behavior analysis
- The intervention targets the core deficits of an autism spectrum disorder, as outlined by the *Diagnostic and Statistical Manual of Mental Disorders*, 5th edition (DSM-5®)
- The intervention is delivered in a home or community setting
- The intervention is rendered directly by a Board-certified Behavior Analyst (BCBA), a licensed mental health clinician with additional documented training in applied behavior analysis, or a paraprofessional under the direct supervision of such professionals
- The intervention is delivered with an appropriate level of intensity (e.g., per Behavior Analyst Certification Board® practice guidelines) and includes ongoing measurement of efficacy: the use of measurement tools and analysis of progress should be continuous, and treatment decisions based on objective analysis of assessment results

Many states have mandated coverage for treatment of autism spectrum disorder.

<http://www.asha.org/Advocacy/state/States-specific-Autism-Mandates/>

Applied Behavior Analysis is unproven for any of the following:

- Programs or interventions that do not meet all of the above proven conditions
- Programs that are not delivered by or under the supervision of an ABA-trained professional
- Programs that target mental disorders other than autism spectrum disorders as defined in the DSM-5®

Services that are otherwise covered under the Individuals with Disabilities Education Act (IDEA) are not covered (e.g., a 1:1 aid in the school setting). School ABA services do allow for coordination of services and would cover services such as, teacher training, meetings with school personnel, and observations in the school setting.

According to a number of recent systematic reviews and meta-analyses, early intervention based on applied behavior analysis is associated with positive outcomes for children with autism spectrum disorder. Currently, there is insufficient evidence to determine which children are most likely to benefit (or not benefit) from specific interventions. Recent progress has been made in systematizing intervention approaches and measuring treatment fidelity.

Intervention research for adolescent and young adult populations and for primary diagnoses other than autism spectrum disorder remains very limited.

UTILIZATION MANAGEMENT CRITERIA

Prior authorization is required for applied behavior analysis (ABA).

Diagnostic Evaluation

The diagnosis of autism spectrum disorder (ASD) must be validated by a documented comprehensive assessment demonstrating the presence of the following diagnostic criteria based on the DSM-5®:

Persistent deficits in social communication and social interaction across multiple contexts, as manifested by the following examples, currently or by history:

- Deficits in social-emotional reciprocity, ranging from abnormal social approach and failure of normal back-and-forth conversation; to reduced sharing of interests, emotions, or affect; to failure to initiate or respond to social interactions.
- Deficits in nonverbal communicative behaviors used for social interaction, ranging from poorly integrated verbal and nonverbal communication; to abnormalities in eye contact and body language or deficits in understanding and use of gestures; to a total lack of facial expressions and non-verbal communication.
- Deficits in developing, maintaining, and understanding relationships, ranging from difficulties adjusting behavior to suit various social contexts; to difficulties in sharing imaginative play or in making friends; to absence of interest in peers.
- Symptoms that impair function are required to in order to be diagnosed with ASD (Hyman et al., 2020).

Specify current severity. See Appendix A.

Restricted, repetitive patterns of behavior, interests, or activities, as manifested by at least two of the following examples, currently or by history:

- Stereotyped or repetitive motor movements, use of objects or speech (e.g., simple motor stereotypes, lining up toys or flipping objects, echolalia, idiosyncratic phrases).
- Insistence on sameness, inflexible adherence to routines, or ritualized patterns of verbal or nonverbal behavior (e.g., extreme distress at small changes, difficulties with transitions, rigid thinking patterns, greeting rituals, need to take same route or eat same food every day).
- Highly restricted, fixated interests that are abnormal in intensity or focus (e.g., strong attachment to or preoccupation with unusual objects, excessively circumscribed or perseverative interests).
- Hyper- or hyporeactivity to sensory input or unusual interest in sensory aspects of the environment (e.g., apparent indifference to pain/temperature, adverse response to specific sounds or textures, excessive smelling or touching of objects, visual fascination with lights or movement).

Specify current severity. See Appendix A.

Symptoms must be present in the early developmental period (but may not become fully manifested until social demands exceed limited capacities, or may be masked by learned strategies in later life).

Symptoms cause clinically significant impairment in social, occupational, or other important areas of current functioning.

These disturbances are not better explained by intellectual disability (intellectual developmental disorder) or global developmental delay. Intellectual disability and autism spectrum disorder frequently co-occur; to make comorbid diagnoses of autism spectrum disorder and intellectual disability, social communication should be below that expected for general developmental level.

Intellectual disability without Autism may be difficult to differentiate from Autism in very young children. Individuals with intellectual disability who have not developed language or symbolic skills also present a challenge for differential diagnosis, since repetitive behavior often occurs in such individuals as well. A diagnosis of Autism in an individual with intellectual disability is appropriate when social communication and interaction are significantly impaired relative to the developmental level of the individual's nonverbal skills (e.g., fine motor skills, nonverbal problem solving). In contrast, intellectual disability is the appropriate diagnosis when there is no apparent discrepancy between the level of social communicative skills and other intellectual skills (DSM-5, Differential Diagnosis, Autism Spectrum Disorder). As clinically indicated, the autism evaluation should include (Volkmar et al., 2014; Meyers & Johnson, 2007, reaffirmed 2014):

- The use of a standard parent- or clinician-rated screening instrument for Autism, such as (Volkmar et al., 2014):
 - Autism Behavior Checklist [ABC]
 - Childhood Autism Rating Scale [CARS]

- Checklist for Autism in Toddlers [CHAT; M-CHAT]
 - Communication and Symbolic Behavior Scales Developmental Profile Infant-Toddler Checklist [CSBS-DP-IT-Checklist]
 - Autism Screening Questionnaire [ASQ]
 - Autism Quotient [AQ]
 - Childhood Autism Screening Test [CAST]
- An essential component of the diagnostic process is formal examinations of language, cognitive, and adaptive abilities and sensory status (Hyman et al., 2020).
- The use of a standard psychiatric assessment for Autism, such as (Volkmar et al., 2014):
 - Autism Diagnostic Interview-Revised [ADI]
 - Autism Diagnostic Observation Schedule [ADOS]
 - Diagnostic Interview for Social and Communication Disorders [DISCO].
- Observation tools used to confirm the ASD diagnosis include the Autism Diagnostic Observation Schedule, Second Edition (ADOS-2) and the Childhood Autism Rating Scale, Second Edition (CARS-2) (Hyman et al., 2020).
- Interviews with the child and family, and assessment of the parents' knowledge of autism spectrum disorder, coping skills, and available resources and supports (Volkmar et al., 2014).
- Review of past records (e.g., past and current behavioral interventions) and historical information (e.g., family history and relevant psychosocial issues) (Volkmar et al., 2014).
- A thorough history includes a long-term experience with the individual that demonstrates the effects of symptoms on the individual's ability to function various settings such as family, peer, and school (Hyman et al., 2020).
 - Questionnaires that establish a history of ASD symptoms and may be used as part of the complete evaluation are the Social Communication Questionnaire (SCQ) or the Social Responsiveness Scale (SRS) (Hyman et al., 2020).
- As clinically appropriate, systematic attention to the areas relevant to differential diagnosis with specific attention to as to whether social communication skills fall below the individual's developmental level, including (*Diagnostic and Statistical Manual of Mental Disorders* 5th ed.; DSM-5; APA, 2013):
 - Rett syndrome
 - Selective mutism
 - Language disorders and social (pragmatic) communication disorder
 - Intellectual disability (intellectual developmental disorder) without autism spectrum disorder
 - Stereotypic movement disorder
 - Attention-deficit/hyperactivity disorder
 - Schizophrenia
- Assessment of co-occurring developmental conditions should include (Hyman et al., 2020):
 - Cognitive Testing
 - Adaptive Function Testing
 - Sensory Assessments: Hearing, Vision, Sensory Processing
- As clinically appropriate, attention to possible comorbid diagnoses
- Observation of broad areas of social interaction and restricted, repetitive behaviors
- When clinically appropriate, a medical assessment, including physical examination, hearing screen, and examination for signs of other genetic abnormalities (Volkmar et al., 2014):
- Identifying the genetic aspect of ASD via genetic testing provides clinicians with additional data for families about prognosis and recurrence risk (Hyman et al., 2020).
- When clinically appropriate, psychological assessment, such as:
 - Measurements of cognitive ability and adaptive skills
 - Use of standard tests of intelligence
 - Identification of areas of strength and weakness useful for designing intervention programs
- When clinically appropriate communication assessment, such as measurement of receptive and expressive vocabulary and language use or a summary of the individual's

- use of language in everyday situations (Volkmar et al., 2014).
- When members of multiple disciplines are involved in assessment (e.g., occupational therapy, physical therapy), coordination among the various professionals is required (Volkmar et al., 2014).

Treatment Planning

Once an ASD diagnosis has been established:

- A standardized functional assessment is used to maximize the effectiveness and efficiency of behavioral support interventions (Myers & Johnson, reaffirmed 2014). The assessment may incorporate information such as interviews with caregivers, structured rating scales, direct observation data, and attention to coexisting medical conditions. The assessment should include baseline data and inform subsequent establishment of treatment goals (Behavior Analyst Certification Board [BACB], 2014).
- When an individual displays maladaptive behaviors it is recommended the credentialed provider complete a functional behavior assessment to better inform treatment planning.
- Targets include areas such as the following:
 - Social communication skills and focus on the social importance of the behaviors targeted
 - Social language skills
 - Social interaction skills
 - Restricted, repetitive patterns of behavior, interests, or activities
 - Self-injurious, violent, destructive or other maladaptive behavior
- A credentialed provider with ABA expertise is identified to provide treatment. Examples include (BACB, 2014):
 - A Master- or Doctoral-level provider that is a Board Certified Behavior Analyst (BCBA)
 - A licensed behavioral health clinician who has attested to having sufficient expertise and has been credentialed to provide ABA services
 - A Board Certified Assistant Behavior Analyst (BCaBA) or non-licensed individual under the direct supervision of a BCBA or licensed behavioral health clinician who takes responsibility for the member's care that does either of the following:
 - Assist in the initial or concurrent assessment of the member's deficits or adaptive behaviors
 - Implement a treatment plan that has been developed by a BCBA or licensed behavioral health clinician
 - Supervision is responsive to individual client needs. Two hours for every ten hours of direct treatment is the general standard of care (BACB, 2014).
 - Direct supervision time may account for 50 percent or more of case supervision time, with the remaining time utilized in indirect supervisory activities such as treatment planning (BACB, 2014).
- Outcome-oriented interventions targeting specific baseline behaviors are identified in a treatment plan describing the frequency, intensity, duration and progress that will be continuously updated.
 - Treatment planning is considered a necessary part of ongoing ABA treatment and should be completed as clinical indicated.
 - The treatment plan must address how the parents/guardians will be trained in management skills that can be generalized to the home.
 - As clinically indicated, parent/guardian training is an expectation. In the rare circumstance that parent/guardian is unable the documentation must reflect the reason and identify an alternate plan to provide management skills in the home.
 - The treatment goals and objectives must be comprehensive and clearly stated.
 - Direct support and training of family members and other professionals promotes optimal functioning and generalization and maintenance of behavioral improvements (BACB, 2014).
 - The treatment plan is coordinated with other professionals to ensure appropriate client progress this may include coordination with the school and applicable IFSP/IEP, outpatient behavioral clinicians, medical doctors, speech/occupational therapists and others (BACB, 2014).

- All components of the child’s care are tracked and updated throughout the duration of services.

Treatment

ABA intervention must include the following elements (BACB, 2014; Myers & Johnson, 2014; Volkmar et al., 2014):

- Mitigate the core features of ASD
- ABA is an intensive treatment
- Target specific deficits related to imitation, attention, motivation, compliance and initiation of interaction, and the specific behaviors that are to be incrementally taught and positively reinforced tie to objective and quantifiable treatment goals that have baseline data, measurable progress, and projected timeframes for completion. Include the child’s parents in parent training and the acquisition of skills in behavior modification to promote management of skills within the home
- Treatment goals are prioritized in to address behaviors that threaten the health or safety of the client or others or create a barrier to quality of life. Goals are also prioritized to increase skills fundamental to maintaining health and social inclusion
- Train family members and other caregivers to manage problem behavior and interact with the child in a therapeutic manner
- As indicated, include psychotherapy (e.g., cognitive behavioral therapy) for higher functioning children to treat conditions such as anxiety and anger management
- Have an appropriate level of intensity and duration driven by factors such as:
 - Treatment plan should indicate the treatment setting, instructional methods to be used, hours requested and clinical justification of those hours
 - Treatment goals that relate to and include how skills will be generalized and maintained across people and environments
 - Changes in the targeted behavior(s) / response to treatment
 - The demonstration and maintenance of management skills by the parents and caregivers;
 - Whether specific issues are being treated in a less intensive group format (e.g., social skills groups)
 - The child’s ability to participate in ABA given attendance at school, daycare or other treatment settings
 - The impact of co-occurring behavioral or medical conditions on skill attainment
 - The member’s overall symptom severity; and
 - The member’s progress in treatment related to treatment duration.

Treatment methodologies utilized as part of intensive behavior therapies should be considered established by the National Autism Centers Standards Projects.

Parent/Caregiver support is expected to be a component of the ABA program, as they will need to provide additional hours of behavioral interventions. Parents or caregivers must be involved and engaged in the training and follow through on treatment recommendations beyond that provided by licensed or certified practitioners. Parent support groups are considered not medically necessary (Myers & Johnson, 2014).

Parent and caregiver training include a systematic, individualized curriculum on ABA fundamental concepts. The goal of this training is skills development and support so that parents and caregivers are proficient in implementing treatment strategies in a variety of settings and critical environments (BACB, 2014).

ABA programs typically fall into either focused or comprehensive ABA treatment. The type of treatment may lend itself to different intensity of services. Total intensity of services includes both direct and indirect services (e.g. caregiver training and supervision). Hours may be increased or decreased based on the client’s response to treatment and current needs (BACB, 2014).

Coordination of Care

If applicable, documentation of communication and coordination with other service providers and agencies, (i.e. day care, preschool, school, early intervention services providers) and/or other allied health care providers (i.e. occupational therapy, speech therapy, physical therapy and any other applicable providers) to reduce the likelihood of unnecessary duplication of services. According to the BACB (2014), collaborating between all professionals engaged with a child will ensure consistency, as better consistency likely leads to better outcomes. Documentation should include the following:

- Types of therapy provided
- Number of therapies per week
- Behaviors/deficits targeted
- Progress related to the treatment/services being provided
- Measureable criteria for completing treatment with projected plan for continued care after discharge from ABA therapy
- Total number of days per week and hours per day of direct services to child and parents or caregivers to include duration and location of requested ABA therapy
- Dates of service requested
- Licensure, certification and credentials of the professionals providing ABA services to the child
- Evidence that parents and/or caregivers have remained engaged in the treatment plan, following all appropriate treatment recommendations
 - Detailed description of interventions with the parent(s) or caregiver(s), including:
 - Parental or caregiver education, training, coaching and support
 - Overall parent or caregiver goals including a brief summary of progress. As part of the summary of progress the information should also include percentage of planned sessions attended
 - Plan for transitioning ABA interventions identified for the child to the parents or caregivers.

Continued treatment

With each medical necessity review for continued ABA treatment, an updated treatment plan and progress reports will be required for review, including all of the following documentation (BACB, 2014; Myers & Johnson, 2014; Volkmar et al., 2014):

- There is a reasonable expectation on the part of the treating clinician that the child's behavior and skill deficits will continue to improve to a clinically meaningful extent, in at least two settings (home, school, community) with ABA services;
- Therapy is not making the symptoms or behaviors persistently worse;
- Progress is assessed and documented for each targeted symptom and behavior, including progress toward defined goals, and including the same modes of measurement that were utilized for baseline measurements of specific symptoms and behaviors;
- The treatment plan and progress report should reflect improvement from baseline in skill deficits and problematic behaviors using validated assessments of adaptive functioning;
- Parent/Caregivers are involved and making progress in their own development of behavioral interventions;
- The treatment plan should reflect a plan to transition services in intensity over time;
- When there has been inadequate progress with targeted symptoms or behaviors, or no demonstrable progress within a six month period, or specific goals have not been achieved within the estimated timeframes, there should be an assessment of the reasons for inadequate progress or not meeting the goals, and treatment interventions should be modified or changed in order to attempt to achieve adequate progress. Documentation of such an assessment and subsequent treatment plan change(s) must include:
 - Increased time and/or frequency working on targets
 - Change in treatment techniques
 - Increased parent/caregiver training
 - Identification and resolution of barriers to treatment effectiveness
 - Any *newly identified* co-existing disorder (e.g., anxiety, psychotic disorder, mood disorder)
 - Goals reconsidered (e.g., modified or removed)

When goals have been achieved, either new goals should be identified that are based on targeted symptoms and behaviors that are preventing the child from adequately participating in age-appropriate home, school or community activities, or that are presenting a safety risk to self, others, or property; or, the treatment plan should be revised to include a transition to less intensive interventions.

Treatment methodologies utilized as part of intensive behavior therapies should be considered established by the National Autism Centers Standards Projects.

Discharge

When any of the following criteria are met the child will be considered discharged and any further ABA services will be considered not medically necessary:

- Documentation that the child demonstrates improvement from baseline in targeted skill deficits and behaviors to the extent that goals are achieved or maximum benefit has been reached
- Documentation that there has been no clinically significant progress or measurable improvement for a period of at least 3 months in the child's behaviors or skill deficits in any of the following measures:
 - Adaptive functioning
 - Communication skills
 - Language skills
 - Social skills
- The treatment is making the skill deficits and/or behaviors persistently worse
- The child is unlikely to continue to benefit or maintain long term gains from continued ABA therapy
- Parents and/or caregivers have refused treatment recommendations or are unable to participate in the treatment program and/or do not follow through on treatment recommendations to an extent that compromises the effectiveness of the services.

Documentation Requirements

ABA providers are required to have a separate record for each member that contains the following documentation:

- Comprehensive assessment establishing the autism diagnosis
- All necessary demographic information
- Complete developmental history and educational assessment
- Functional behavioral assessment including assessment of targeted risk behaviors
- Behavioral/medical health treatment history including but not limited to:
 - known conditions
 - dates and providers of previous treatment
 - current treating clinicians
 - current therapeutic interventions and responses
- Individualized treatment plan and all revisions to the treatment plan, including objective and measurable goals, as well as parent training
- Daily progress notes including:
 - place of service
 - start and stop time
 - who rendered the service
 - the specific service (e.g., parenting training, supervision, direct service)
 - who attended the session
 - interventions that occurred during the session
 - barriers to progress
 - response to interventions
- All documentation must be legible
- All documentation related to coordination of care
- All documentation related to supervision of paraprofessionals
- If applicable, a copy of the child's Individualized Education Plan (IEP)

- If applicable, progress notes related to Early Intervention Plan or Pre-school/Special Education Program or allied health services
- Certification and credentials of the professionals providing the ABA therapy.

APPENDIX A

Severity Levels for Autism Spectrum Disorder

Severity Level	Social Communication	Restricted, Repetitive Behaviors
Level 3 –Requiring very substantial support	<i>Severe deficits in verbal and nonverbal social communication skills causes severe impairments in functioning, very limited initiation of social interactions, and minimal response to social overtures from others. For example, a person with few words of intelligible speech who rarely initiates interaction and, when he or she does, makes unusual approaches to meet needs only and responds to only very direct social approaches.</i>	<i>Inflexibility of behavior, extreme difficulty coping with change, or other restricted/repetitive behaviors markedly interferes with functioning in all spheres. Great distress/difficulty in changing focus or action.</i>
Level 2 –Requiring substantial support	<i>Marked deficits in verbal and nonverbal social communication skills; social impairments apparent even with supports in place; limited initiation of social interactions; and reduced or abnormal responses to social overtures from others. For example, a person who speaks simple sentences, whose interaction is limited to narrow special interests, and who has markedly odd nonverbal communication.</i>	<i>Inflexibility of behavior, difficulty coping with change, or other restricted/repetitive behaviors appear frequently enough to be obvious to the casual observer and interfere with functioning in a variety of contexts. Distress and/or difficulty changing focus or action.</i>
Level 1 –Requiring support	<i>Without supports in place, deficits in social communication cause noticeable impairments. Difficulty initiating social interactions, and clear examples of atypical or unsuccessful responses to social overtures of others. May appear to have decreased interest in social interactions. For example, a person who is able to speak in full sentences and engages in communication but whose to-and-fro conversation with others fails, and whose attempts to make friends are odd and typically unsuccessful.</i>	<i>Inflexibility of behavior causes significant interference with functioning in or more contexts. Difficulty switching between activities. Problems of organization and planning hamper independence.</i>

Source: American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*, Fifth edition. Table 2. American Psychiatric Publishing.

APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member-specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other clinical criteria may apply.

Diagnosis Codes	Description
F84.0	Autistic Disorder

CPT Code	Description
97151	Behavior identification assessment, administered by a physician or other qualified healthcare professional, each 15 minutes of the physician's or other qualified healthcare professional's time face-to-face with patient and/or guardian(s)/caregiver(s) administering assessments and discussing findings and recommendations, and non-face-to-face analyzing past data, scoring/interpreting the assessment, and preparing the report/treatment plan.
97152	Behavior identification supporting assessment, administered by one technician under the direction of a physician or other qualified healthcare professional, face-to-face with the patient, each 15 minutes.
97153	Adaptive behavior treatment by protocol, administered by a technician under the direction of a physician or other qualified healthcare professional, face-to-face with one patient, every 15 minutes.
97154	Group adaptive behavior treatment by protocol, administered by a technician under the direction of a physician or other qualified healthcare professional, with two or more patients, every 15 minutes.
97155	Adaptive behavior treatment with protocol modification, administered by a physician or other qualified healthcare professional, which may include simultaneous direction of a technician, face-to-face with one patient, every 15 minutes.
97156	Family adaptive behavior treatment guidance, administered by a physician or other qualified healthcare professional (with or without the patient present), face-to-face with guardian(s)/caregiver(s), every 15 minutes.
97157	Multiple-family group adaptive behavior treatment guidance, administered by a physician or other qualified healthcare professional (without the patient present), face-to-face with multiple sets of guardians/caregivers, every 15 minutes.
97158	Group adaptive behavior treatment with protocol modification, administered by a physician or other qualified healthcare professional, face-to-face with multiple patients, every 15 minutes.
0362T	Behavior identification supporting assessment, every 15 minutes of technicians' time face-to-face with a patient, requiring the following components: <ul style="list-style-type: none"> •administered by the physician or other qualified healthcare professional who is on site; •with the assistance of two or more technicians; •for a patient who exhibits destructive behavior; •completed in an environment that is customized to the patient's behavior.
0373T	Adaptive behavior treatment with protocol modification, every 15 minutes of technicians' time face-to-face with a patient, requiring the following components: <ul style="list-style-type: none"> • administered by the physician or other qualified healthcare professional who is on site; •with the assistance of two or more technicians; •for a patient who exhibits destructive behavior; •completed in an environment that is customized, to the patient's behavior.

Procedure Codes	Description
H0031	Mental health assessment, by non-physician
H0032	Mental health service plan development, by non-physician
H2012	Behavioral health day treatment, per hour
H2014	Skills training and development, per 15 minutes

H2019	Therapeutic behavioral services, per 15 minutes
H2021 (Pennsylvania only)	Community-based wrap around services, per 15 minutes
H2027 (Pennsylvania only)	Psychoeducational service, per 15 minutes

REFERENCES

1. American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders*, Fifth Edition. American Psychiatric Publishing.
2. Behavior Analyst Certification Board. (2014). *Applied Behavior Analysis Treatment of Autism Spectrum Disorder: Practice Guidelines for Healthcare Funders and Managers*. Second edition. Copyright © by the Behavior Analyst Certification Board, Inc. (BACB), all rights reserved.
3. Hyman, S.L., Levy, S.E., & Myers, S.M. and the American Academy of Pediatrics Council on Children With Disabilities, Section On Developmental And Behavioral Pediatrics. (2020). Identification, evaluation, and management of children with autism spectrum disorder. *Pediatrics*, 145(1), 1-71.
4. Myers, S.M. & Johnson, C.P. and the American Academy of Pediatrics Council on Children with Disabilities. (2007, reaffirmed 2014). Management of children with autism spectrum disorders. *Pediatrics*, 120(5),1162-1182.
5. Volkmar, F., Siegel, M., Woodbury-Smith, M., King, B., McCracken, J., State, M. and the American Academy of Child and Adolescent Psychiatry (AACAP) Committee on Quality Issues (CQI). (2014). Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 53(2), 237-257.

REVISION HISTORY

Date	Action/Description
03/16/2020	<ul style="list-style-type: none"> • Version 1; Supplemental Clinical Criteria